

Market Applicability													
Market	DC	FL & FHK	FL MMA	FL LTC	GA	KY	MD	NJ	NV	NY	TN	TX	WA
Applicable	X	X	NA	NA	X	X	X	X	X	X	NA	NA	NA

*FHK- Florida Healthy Kids

Hyrimoz (adalimumab-adaz)

Override(s)	Approval Duration
Prior Authorization Quantity Limit	1 year

Medications	Quantity Limit
Hyrimoz (adalimumab-adaz) 40 mg/0.8 mL prefilled pen/syringe ^{#* ‡^}	2 pens/syringes per 28 days

Override Criteria

[#]In the treatment of Rheumatoid Arthritis (RA): May approve up to 4 (four) syringes, autoinjectors, or pens (40mg/0.8 mL) [up to an additional 2 (two) syringes, autoinjectors, or pens] every 28 days if the individual is unable to take concomitant methotrexate.

^{*}Initiation of therapy for adult Crohn's Disease (CD) or Ulcerative Colitis (UC): May approve up to 4 (four) additional pens, autoinjectors, or syringes (40 mg/0.8 mL) in the first month (28 days) of treatment.

[‡]In the treatment of CD or UC: May approve up to an additional 2 (two) syringes, autoinjectors, or pens (40 mg/0.8 mL) every 28 days if the individual has an inadequate response to standard maintenance dosing.

[^]Initiation of therapy for Plaque Psoriasis (Ps) (Psoriasis vulgaris): May approve up to 2 (two) additional pens, autoinjectors, or syringes (40 mg/0.8 mL) in the first month (28 days) of treatment.

APPROVAL CRITERIA

Requests for Hyrimoz (adalimumab-adaz) may be approved for the following:

- I. Crohn's disease (CD) when each of the following criteria are met:
 - A. Individual is 6 years of age or older with moderate to severe CD;

AND

 - B. Individual has had an inadequate response to, is intolerant of, or has a contraindication to conventional therapy (such as 5-Aminosalicylic acid products, systemic corticosteroids, or immunosuppressants);

PAGE 1 of 11 05/22/2019
New Program Date 03/01/2019

This policy does not apply to health plans or member categories that do not have pharmacy benefits, nor does it apply to Medicare. Note that market specific restrictions or transition-of-care benefit limitations may apply.

CRX-ALL-0392-19

Market Applicability													
Market	DC	FL & FHK	FL MMA	FL LTC	GA	KY	MD	NJ	NV	NY	TN	TX	WA
Applicable	X	X	NA	NA	X	X	X	X	X	X	NA	NA	NA

*FHK- Florida Healthy Kids

AND

- C. Individual has had a trial (medication samples/coupons/discount cards are excluded from consideration as a trial) and inadequate response or intolerance to ONE (1) preferred biologic agent [Current preferred biologics include - Humira (adalimumab), Inflectra (infliximab-dyyb), Renflexis (infliximab-abda)] unless the following criteria are met:
1. Individual has been receiving and is maintained on a stable dose of Hyrimoz (adalimumab-adaz); **OR**
 2. The preferred agent is not FDA-approved and do not have an accepted off-label use per the off-label policy for the prescribed indication and Hyrimoz (adalimumab-adaz) does; **OR**
 3. The preferred agent is not acceptable due to concomitant clinical conditions, such as but not limited to any of the following:
 - a. Known hypersensitivity to any active or inactive component which is not also associated with Hyrimoz (adalimumab-adaz); **OR**
 - b. Individual's age; **OR**
 - c. Pregnant or planning on becoming pregnant; **OR**
 - d. Serious infections or concurrent sepsis; **OR**
 4. The preferred agent(s) do not have activity against a concomitant clinical condition and Hyrimoz (adalimumab-adaz) does. Examples include but may not be limited to the following:
 - a. Concomitant Psoriasis: TNFi (agents FDA-approved for both indications) or Stelara are preferred; **OR**
 - b. Concomitant Rheumatoid Arthritis: TNFi agents are preferred;

OR

- II. Ulcerative colitis (UC) when each of the following criteria are met:
- A. Individual is 18 years of age or older with moderate to severe UC;

AND

- B. Individual has had an inadequate response to, is intolerant of, or has a contraindication to conventional therapy (such as 5-Aminosalicylic acid products, systemic corticosteroids, or immunosuppressants);

AND

- C. Individual has had a trial (medication samples/coupons/discount cards are excluded from consideration as a trial) and inadequate response or intolerance to ONE (1) preferred biologic agent [Current preferred biologics include - Humira (adalimumab), Inflectra (infliximab-dyyb), Renflexis (infliximab-abda)] unless the following criteria are met:

This policy does not apply to health plans or member categories that do not have pharmacy benefits, nor does it apply to Medicare. Note that market specific restrictions or transition-of-care benefit limitations may apply.

Market Applicability													
Market	DC	FL & FHK	FL MMA	FL LTC	GA	KY	MD	NJ	NV	NY	TN	TX	WA
Applicable	X	X	NA	NA	X	X	X	X	X	X	NA	NA	NA

*FHK- Florida Healthy Kids

1. Individual has been receiving and is maintained on a stable dose of Hyrimoz (adalimumab-adaz); **OR**
2. The preferred agent is not FDA-approved and do not have an accepted off-label use per the off-label policy for the prescribed indication and Hyrimoz (adalimumab-adaz) does; **OR**
3. The preferred agent is not acceptable due to concomitant clinical conditions, such as but not limited to any of the following:
 - a. Known hypersensitivity to any active or inactive component which is not also associated with Hyrimoz (adalimumab-adaz); **OR**
 - b. Individual's age; **OR**
 - c. Pregnant or planning on becoming pregnant; **OR**
 - d. Serious infections or concurrent sepsis; **OR**
4. The preferred agent(s) do not have activity against a concomitant clinical condition and Hyrimoz (adalimumab-adaz) does. Examples include but may not be limited to the following:
 - a. Concomitant Psoriasis: TNFi (agents FDA-approved for both indications) or Stelara are preferred; **OR**
 - b. Concomitant Rheumatoid Arthritis: TNFi agents are preferred.

OR

- III. Rheumatoid arthritis (RA) when each of the following criteria are met:
 - A. Individual must be 18 years of age or older with moderate to severe RA;

AND

 - B. Individual has had an inadequate response to, is intolerant of, or has a contraindication to conventional therapy [nonbiologic disease modifying anti-rheumatic agents (DMARDs) (such as methotrexate, sulfasalazine, leflunomide, or hydroxychloroquine)] (ACR 2015);

AND

 - C. Individual has had a trial (medication samples/coupons/discount cards are excluded from consideration as a trial) and inadequate response or intolerance to TWO (2) preferred biologic agents [Current preferred biologics include – Enbrel (etanercept), Humira (adalimumab)] unless the following criteria are met:
 1. Individual has been receiving and is maintained on a stable dose of Hyrimoz (adalimumab-adaz); **OR**
 2. The preferred agents are not acceptable due to concomitant clinical conditions, such as but not limited to any of the following:
 - a. Known hypersensitivity to any active or inactive component which is not also associated with Hyrimoz (adalimumab-adaz); **OR**
 - b. Individual's age; **OR**

This policy does not apply to health plans or member categories that do not have pharmacy benefits, nor does it apply to Medicare. Note that market specific restrictions or transition-of-care benefit limitations may apply.

Market Applicability													
Market	DC	FL & FHK	FL MMA	FL LTC	GA	KY	MD	NJ	NV	NY	TN	TX	WA
Applicable	X	X	NA	NA	X	X	X	X	X	X	NA	NA	NA

*FHK- Florida Healthy Kids

- c. Pregnant or planning on becoming pregnant; **OR**
- d. Serious infections or concurrent sepsis; **OR**
- 3. The preferred agent(s) do not have activity against a concomitant clinical condition and Hyrimoz (adalimumab-adaz) does. An example includes but may not be limited to the following:
 - a. Concomitant Crohn's disease: TNFi (agents FDA-approved for both indications) are preferred;

OR

- IV. Ankylosing spondylitis (AS) when each of the following criteria are met:
 - A. Individual is 18 years of age or older with moderate to severe AS;
 - AND**
 - B. Individual has had an inadequate response to, is intolerant of, or has a contraindication to conventional therapy [such as NSAIDs or nonbiologic disease modifying anti-rheumatic agents (DMARDs) (such as sulfasalazine)];
 - AND**
 - C. Individual has had a trial (medication samples/coupons/discount cards are excluded from consideration as a trial) and inadequate response or intolerance to TWO (2) preferred biologic agents [Current preferred biologics include – Enbrel (etanercept), Humira (adalimumab)] unless the following criteria are met:
 - 1. Individual has been receiving and is maintained on a stable dose of Hyrimoz (adalimumab-adaz); **OR**
 - 2. The preferred agents are not FDA-approved and do not have an accepted off-label use per the off-label policy for the prescribed indication and Hyrimoz (adalimumab-adaz) does; **OR**
 - 3. The preferred agents are not acceptable due to concomitant clinical conditions, such as but not limited to any of the following:
 - a. Known hypersensitivity to any active or inactive component which is not also associated with Hyrimoz (adalimumab-adaz); **OR**
 - b. Individual's age; **OR**
 - c. Pregnant or planning on becoming pregnant; **OR**
 - d. Serious infections or concurrent sepsis;

OR

- V. Polyarticular juvenile idiopathic arthritis (PJIA) when each of the following criteria are met:
 - A. Individual is 2 years of age or older with moderate to severe(PJIA);
 - AND**

This policy does not apply to health plans or member categories that do not have pharmacy benefits, nor does it apply to Medicare. Note that market specific restrictions or transition-of-care benefit limitations may apply.

Market Applicability													
Market	DC	FL & FHK	FL MMA	FL LTC	GA	KY	MD	NJ	NV	NY	TN	TX	WA
Applicable	X	X	NA	NA	X	X	X	X	X	X	NA	NA	NA

*FHK- Florida Healthy Kids

B. Individual has had an inadequate response to, is intolerant of, or has a contraindication to conventional therapy [nonbiologic disease modifying anti-rheumatic drugs (DMARDs) (such as methotrexate, sulfasalazine, or leflunomide)];

AND

C. Individual has had a trial (medication samples/coupons/discount cards are excluded from consideration as a trial) and inadequate response or intolerance to TWO (2) preferred biologic agents [Current preferred biologics include – Enbrel (etanercept), Humira (adalimumab)] unless the following criteria are met:

1. Individual has been receiving and is maintained on a stable dose of Hyrimoz (adalimumab-adaz); **OR**
2. The preferred agents are not FDA-approved and do not have an accepted off-label use per the off-label policy for the prescribed indication and Hyrimoz (adalimumab-adaz) does; **OR**
3. The preferred agents are not acceptable due to concomitant clinical conditions, such as but not limited to any of the following:
 - a. Known hypersensitivity to any active or inactive component which is not also associated with Hyrimoz (adalimumab-adaz); **OR**
 - b. Individual's age; **OR**
 - c. Pregnant or planning on becoming pregnant; **OR**
 - d. Serious infections or concurrent sepsis;

OR

VI. Psoriatic arthritis (PsA) when each of the following criteria are met:

A. Individual is 18 years of age or older with moderate to severe PsA;

AND

B. Individual has had an inadequate response to, is intolerant of, or has had a contraindication to conventional therapy [nonbiologic disease modifying anti-rheumatic drugs (DMARDs) (such as methotrexate, sulfasalazine, or leflunomide)];

AND

C. Individual has had a trial (medication samples/coupons/discount cards are excluded from consideration as a trial) and inadequate response or intolerance to TWO (2) preferred biologic agents [Current preferred biologics include – Enbrel (etanercept), Humira (adalimumab)] unless the following criteria are met:

1. Individual has been receiving and is maintained on a stable dose of Hyrimoz (adalimumab-adaz); **OR**
2. The preferred agents are not FDA-approved and do not have an accepted off-label use per the off-label policy for the prescribed indication and Hyrimoz (adalimumab-adaz) does; **OR**

This policy does not apply to health plans or member categories that do not have pharmacy benefits, nor does it apply to Medicare. Note that market specific restrictions or transition-of-care benefit limitations may apply.

Market Applicability													
Market	DC	FL & FHK	FL MMA	FL LTC	GA	KY	MD	NJ	NV	NY	TN	TX	WA
Applicable	X	X	NA	NA	X	X	X	X	X	X	NA	NA	NA

*FHK- Florida Healthy Kids

3. The preferred agents are not acceptable due to concomitant clinical conditions, such as but not limited to any of the following:
 - a. Known hypersensitivity to any active or inactive component which is not also associated with Hyrimoz (adalimumab-adaz); **OR**
 - b. Individual's age; **OR**
 - c. Pregnant or planning on becoming pregnant; **OR**
 - d. Serious infections or concurrent sepsis; **OR**
4. The preferred agent(s) do not have activity against a concomitant clinical condition and Hyrimoz (adalimumab-adaz) does. Examples include but may not be limited to the following:
 - a. Concomitant Crohn's Disease: TNFi (agents FDA-approved for both indications) or Stelara are preferred; **OR**
 - b. Concomitant Ulcerative Colitis: TNFi (agents FDA-approved for both indications) are preferred;

OR

- VII. Plaque psoriasis (Ps) (Psoriasis vulgaris) when each of the following criteria are met:
- A. Individual is 18 years of age or older with chronic moderate to severe (that is, extensive or disabling) plaque Ps (psoriasis vulgaris) with either of the following (AAD 2011):
 1. Plaque Ps (psoriasis vulgaris) involving greater than five percent (5%) body surface area (BSA); **OR**
 2. Plaque Ps (psoriasis vulgaris) involving less than or equal to five percent (5%) BSA involving sensitive areas or areas that significantly impact daily function (such as palms, soles of feet, head/neck, or genitalia);

AND

- B. Individual has had an inadequate response to, is intolerant of, or has a contraindication to phototherapy or other systemic therapy (such as acitretin, cyclosporine, or methotrexate);

AND

- C. Individual has had a trial (medication samples/coupons/discount cards are excluded from consideration as a trial) and inadequate response or intolerance to TWO (2) preferred biologic agents [Current preferred biologics include – Enbrel (etanercept), Humira (adalimumab)] unless the following criteria are met:
 1. Individual has been receiving and is maintained on a stable dose of Hyrimoz (adalimumab-adaz); **OR**
 2. The preferred agents are not FDA-approved and do not have an accepted off-label use per the off-label policy for the prescribed indication and Hyrimoz (adalimumab-adaz) does; **OR**

This policy does not apply to health plans or member categories that do not have pharmacy benefits, nor does it apply to Medicare. Note that market specific restrictions or transition-of-care benefit limitations may apply.

Market Applicability													
Market	DC	FL & FHK	FL MMA	FL LTC	GA	KY	MD	NJ	NV	NY	TN	TX	WA
Applicable	X	X	NA	NA	X	X	X	X	X	X	NA	NA	NA

*FHK- Florida Healthy Kids

3. The preferred agents are not acceptable due to concomitant clinical conditions, such as but not limited to any of the following:
 - a. Known hypersensitivity to any active or inactive component which is not also associated with Hyrimoz (adalimumab-adaz); **OR**
 - b. Individual's age; **OR**
 - c. Pregnant or planning on becoming pregnant; **OR**
 - d. Serious infections or concurrent sepsis; **OR**
4. The preferred agent(s) do not have activity against a concomitant clinical condition and Hyrimoz (adalimumab-adaz) does. Examples include but may not be limited to the following:
 - a. Concomitant Crohn's Disease: TNFi (agents FDA-approved for both indications) or Stelara are preferred; **OR**
 - b. Concomitant Ulcerative Colitis: TNFi (agents FDA-approved for both indications) are preferred;

OR

- VIII. Non-infectious uveitis (UV) when each of the following criteria are met:
- A. Individual has chronic, recurrent, treatment-refractory or vision-threatening disease;
- AND**
- B. Individual has had an inadequate response to, is intolerant of, or has a contraindication to conventional therapy [such as corticosteroids or immunosuppressants (azathioprine, cyclosporine, or methotrexate)];
- AND**
- C. Individual has had a trial (medication samples/coupons/discount cards are excluded from consideration as a trial) and inadequate response or intolerance to the preferred biologic agent [Current preferred biologic includes – Humira (adalimumab)] unless the following criteria are met:
 1. Individual has been receiving and is maintained on a stable dose of Hyrimoz (adalimumab-adaz); **OR**
 2. The preferred agents are not FDA-approved and do not have an accepted off-label use per the off-label policy for the prescribed indication and Hyrimoz (adalimumab-adaz) does; **OR**
 3. The preferred agents are not acceptable due to concomitant clinical conditions, such as but not limited to any of the following:
 - a. Known hypersensitivity to any active or inactive component which is not also associated with Hyrimoz (adalimumab-adaz); **OR**
 - b. Individual's age; **OR**
 - c. Pregnant or planning on becoming pregnant; **OR**
 - d. Serious infections or concurrent sepsis;

This policy does not apply to health plans or member categories that do not have pharmacy benefits, nor does it apply to Medicare. Note that market specific restrictions or transition-of-care benefit limitations may apply.

Market Applicability													
Market	DC	FL & FHK	FL MMA	FL LTC	GA	KY	MD	NJ	NV	NY	TN	TX	WA
Applicable	X	X	NA	NA	X	X	X	X	X	X	NA	NA	NA

*FHK- Florida Healthy Kids

OR

- IX. Hidradenitis suppurativa (HS) when each of the following criteria are met:
- A. Individual is 12 years of age or older;
- AND**
- B. Individual has moderate to severe hidradenitis suppurativa (Hurley stage II or Hurley stage III disease);
- AND**
- C. Individual has had an inadequate response to, is intolerant of, or has a contraindication to conventional therapy (such as oral antibiotics);
- AND**
- D. Individual has had a trial (medication samples/coupons/discount cards are excluded from consideration as a trial) and inadequate response or intolerance to the preferred biologic agent [Current preferred biologic includes – Humira (adalimumab)] unless the following criteria are met:
 - 1. Individual has been receiving and is maintained on a stable dose of Hyrimoz (adalimumab-adaz); **OR**
 - 2. The preferred agents are not FDA-approved and do not have an accepted off-label use per the off-label policy for the prescribed indication and Hyrimoz (adalimumab-adaz) does; **OR**
 - 3. The preferred agents are not acceptable due to concomitant clinical conditions, such as but not limited to any of the following:
 - a. Known hypersensitivity to any active or inactive component which is not also associated with Hyrimoz (adalimumab-adaz); **OR**
 - b. Individual's age; **OR**
 - c. Pregnant or planning on becoming pregnant; **OR**
 - d. Serious infections or concurrent sepsis.

OR

- X. Sarcoidosis when each of the following criteria are met (Sweiss 2014):
- A. Individual is 18 years of age or older;
- AND**
- B. Individual has chronic, progressive, treatment-refractory disease;
- AND**
- C. Individual has had an inadequate response to, is intolerant of, or has a contraindication to systemic corticosteroids;
- AND**
- D. Individual has had an inadequate response to, is intolerant of, or has a

This policy does not apply to health plans or member categories that do not have pharmacy benefits, nor does it apply to Medicare. Note that market specific restrictions or transition-of-care benefit limitations may apply.

Market Applicability													
Market	DC	FL & FHK	FL MMA	FL LTC	GA	KY	MD	NJ	NV	NY	TN	TX	WA
Applicable	X	X	NA	NA	X	X	X	X	X	X	NA	NA	NA

*FHK- Florida Healthy Kids

contraindication to nonbiologic disease modifying anti-rheumatic agents (DMARDs) (such as methotrexate or azathioprine);

AND

- E. Individual has had a trial (medication samples/coupons/discount cards are excluded from consideration as a trial) and inadequate response or intolerance to the preferred biologic agent [Current preferred biologic includes – Humira (adalimumab)] unless the following criteria are met:
1. Individual has been receiving and is maintained on a stable dose of Hyrimoz (adalimumab-adaz); **OR**
 2. The preferred agents are not FDA-approved and do not have an accepted off-label use per the off-label policy for the prescribed indication and Hyrimoz (adalimumab-adaz) does; **OR**
 3. The preferred agents are not acceptable due to concomitant clinical conditions, such as but not limited to any of the following:
 - a. Known hypersensitivity to any active or inactive component which is not also associated with Hyrimoz (adalimumab-adaz); **OR**
 - b. Individual's age; **OR**
 - c. Pregnant or planning on becoming pregnant; **OR**
 - d. Serious infections or concurrent sepsis.

Hyrimoz (adalimumab-adaz) may **not** be approved for the following:

- I. In combination with other TNF antagonists, JAK inhibitors or other biologic drugs (such as, abatacept, anakinra, or vedolizumab); **OR**
- II. Tuberculosis, other active serious infections, or a history of recurrent infections; **OR**
- III. Individual has not had a tuberculin skin test (TST), or a Centers for Disease Control (CDC-) and Prevention -recommended equivalent, to evaluate for latent tuberculosis prior to initiating adalimumab-adaz.

Note:

TNFi have black box warnings for serious infections and malignancy. Individuals treated with TNFi are at increased risk for developing serious infections that may lead to hospitalization or death. Most individuals who developed these infections were taking concomitant immunosuppressants such as methotrexate or corticosteroids. TNFi should be discontinued if an individual develops a serious infection or sepsis. Individuals should be tested for latent tuberculosis (TB) before TNFi use and during therapy. Treatment for latent TB should be initiated prior to TNFi use. Risks and benefits of TNFi should be carefully considered prior to initiation of therapy in individuals with chronic or recurrent infection. Lymphoma and other

This policy does not apply to health plans or member categories that do not have pharmacy benefits, nor does it apply to Medicare. Note that market specific restrictions or transition-of-care benefit limitations may apply.

Market Applicability													
Market	DC	FL & FHK	FL MMA	FL LTC	GA	KY	MD	NJ	NV	NY	TN	TX	WA
Applicable	X	X	NA	NA	X	X	X	X	X	X	NA	NA	NA

*FHK- Florida Healthy Kids

malignancies have been reported in children and adolescents treated with TNFi. Postmarketing cases of hepatosplenic T-cell lymphoma (HSTCL) have been reported in individuals treated with TNFi. Almost all cases had received treatment with azathioprine or 6-mercaptopurine concomitantly with a TNFi at or prior to diagnosis. It is uncertain whether HSTCL is related to the use of a TNFi or a TNFi in combination with these other immunosuppressants.

State Specific Mandates		
State name	Date effective	Mandate details (including specific bill if applicable)
N/A	N/A	N/A

Key References:

1. Clinical Pharmacology [database online]. Tampa, FL: Gold Standard, Inc.: 2018. URL: <http://www.clinicalpharmacology.com>. Updated periodically.
2. DailyMed. Package inserts. U.S. National Library of Medicine, National Institutes of Health website. <http://dailymed.nlm.nih.gov/dailymed/about.cfm>. Accessed: September 14, 2018.
3. DrugPoints® System [electronic version]. Truven Health Analytics, Greenwood Village, CO. Updated periodically.
4. Lexi-Comp ONLINE™ with AHFS™, Hudson, Ohio: Lexi-Comp, Inc.; 2018; Updated periodically.
5. NCCN Drugs & Biologics Compendium (NCCN Compendium®) 2016 National Comprehensive Cancer Network, Inc. Available at: NCCN.org. Updated periodically. Accessed on: September 14, 2018.
6. Singh JA, Saag KG, Bridges SL et al. 2015 American College of Rheumatology Guideline for the treatment of rheumatoid arthritis. *Arthritis Rheum.* 2016;68:1-26.
7. Menter A, Korman NJ, Elmets CA et al for the American Academy of Dermatology. Guidelines of care for the management of psoriasis and psoriatic arthritis. *J Am Acad Dermatol.* 2011; 65: 137-174.
8. American Gastroenterological Association. Identification, assessment and initial medical treatment of ulcerative colitis Clinical Care Pathway. Available at <https://gastro.org/guidelines/ibd-and-bowel-disorders>. Accessed on: September 14, 2018.
9. American Gastroenterological Association. Identification, assessment and initial medical treatment of Crohn's disease Clinical Care Pathway. Available at <https://gastro.org/guidelines/ibd-and-bowel-disorders>. Accessed on: September 14, 2018.
10. Lichtenstein GR, Loftus EV, Isaacs KL et al. 2018 American College of Gastroenterology Guideline for the management of Crohn's disease in adults. *Am J Gastroenterol* 2018; 113:481–517.
11. Ward MM, Deodhar A, Akl EA, et al. American College of Rheumatology/Spondylitis Association of America/ Spondyloarthritis Research and Treatment Network 2015 Recommendations for the Treatment of Ankylosing Spondylitis and Nonradiographic Axial Spondyloarthritis. *Arthritis Rheumatol.* 2016; 68(2):282-298.
12. Ringold S, Weiss PF, Beukelman T, et al. 2013 Update of the 2011 American College of Rheumatology recommendations for the treatment of juvenile idiopathic arthritis: recommendations for the medical therapy of children with systemic juvenile idiopathic arthritis and tuberculosis screening among children receiving biologic medications. *Arthritis Rheum.* 2013; 65(10):2499-2512.
13. Beukelman T, Patkar NM, Saag KG, et al. 2011 American College of Rheumatology recommendations for the treatment of juvenile idiopathic arthritis: initiation and safety monitoring of therapeutic agents for the treatment of arthritis and systemic features. *Arthritis Care & Research.* 2011; 63(4):465-482.

This policy does not apply to health plans or member categories that do not have pharmacy benefits, nor does it apply to Medicare. Note that market specific restrictions or transition-of-care benefit limitations may apply.

Market Applicability													
Market	DC	FL & FHK	FL MMA	FL LTC	GA	KY	MD	NJ	NV	NY	TN	TX	WA
Applicable	X	X	NA	NA	X	X	X	X	X	X	NA	NA	NA

*FHK- Florida Healthy Kids

14. Levy-Clarke G, Jabs DA, Read RW, et al. Expert panel recommendations for the use of anti-tumor necrosis factor biologic agents in patients with ocular inflammatory disorders; American Uveitis Society subcommittee. Ophthalmology. 2014; 121(3):785-796.
15. Baughman RP, Drent M, et al. Infliximab therapy in patients with chronic sarcoidosis and pulmonary involvement. Am J Respir Crit Care Med. 2006; 174:795-802.
16. Sweiss NJ, Noth I, et al. Efficacy results of a 52-week trial of adalimumab in the treatment of refractory sarcoidosis. Sarcoidosis Vasc Diffuse Lung Dis. 2014; 31(1):46-54.

This policy does not apply to health plans or member categories that do not have pharmacy benefits, nor does it apply to Medicare. Note that market specific restrictions or transition-of-care benefit limitations may apply.