

Market Applicability														
Market	DC	FL & FHK	FL MMA	FL LTC	GA	KS	KY	MD	NJ	NV	NY	TN	TX	WA
Applicable	X	X	NA	NA	X	NA	X	X	X	X	X	NA	NA	NA

*FHK- Florida Healthy Kids

Xeljanz (tofacitinib), Xeljanz XR (tofacitinib extended-release)

Override(s)	Approval Duration
Prior Authorization Quantity Limit	1 year

Medications	Quantity Limit
Xeljanz (tofacitinib)	May be subject to quantity limit
Xeljanz XR (tofacitinib extended-release)	

APPROVAL CRITERIA

Requests for Xeljanz (tofacitinib) or Xeljanz XR (tofacitinib extended-release) may be approved based on the following criteria:

- I. Diagnosis of Rheumatoid Arthritis:
 - A. Individual is 18 years of age or older with moderately to severely active rheumatoid arthritis;

AND

 - B. Agent is used for any of the following reasons:
 1. To reduce signs or symptoms; **OR**
 2. To induce or maintain clinical response; **OR**
 3. To improve physical function;

AND

 - C. Individual has had an inadequate response to, is intolerant of or has a contraindication to methotrexate, sulfasalazine, leflunomide, or hydroxychloroquine);

AND

 - D. Individual has had a trial (medication samples/coupons/discount cards are excluded from consideration as a trial) and inadequate response or intolerance to TWO (2) preferred biologic agents [Current preferred biologics include – Enbrel (etanercept), Humira (adalimumab)] unless the following criteria are met:
 1. Individual has been receiving and is maintained on a stable dose of Xeljanz (tofacitinib) or Xeljanz XR (tofacitinib extended-release); **OR**
 2. The preferred agents are not acceptable due to concomitant clinical conditions, such as but not limited to any of the following:
 - a. Known hypersensitivity to any active or inactive component which is not

PAGE 1 of 5 12/21/2018

This policy does not apply to health plans or member categories that do not have pharmacy benefits, nor does it apply to Medicare. Note that market specific restrictions or transition-of-care benefit limitations may apply.

CRX-ALL-0312-18

Market Applicability														
Market	DC	FL & FHK	FL MMA	FL LTC	GA	KS	KY	MD	NJ	NV	NY	TN	TX	WA
Applicable	X	X	NA	NA	X	NA	X	X	X	X	X	NA	NA	NA

*FHK- Florida Healthy Kids

- also associated with Xeljanz (tofacitinib) or Xeljanz XR (tofacitinib extended-release); **OR**
- b. Individual's age; **OR**
- c. Pregnant or planning on becoming pregnant; **OR**
- d. Serious infections or concurrent sepsis; **OR**
- 3. The individual has either concomitant clinical condition:
 - a. Demyelinating disease; **OR**
 - b. Heart failure with documented left ventricular dysfunction; **OR**
- 4. The preferred agent(s) do not have activity against a concomitant clinical condition and Xeljanz (tofacitinib) or Xeljanz XR (tofacitinib extended-release) does. An example includes but may not be limited to the following:
 - a. Concomitant Crohn's disease: TNFi (agents FDA-approved for both indications) are preferred;

OR

II. Diagnosis of Psoriatic Arthritis:

A. Individual is 18 years of age or older with active psoriatic arthritis;

AND

B. Agent is used for any of the following reasons:

- 1. To reduce signs or symptoms; **OR**
- 2. To induce or maintain clinical response; **OR**
- 3. To improve physical function;

AND

C. Individual has had an inadequate response to, is intolerant of, or has a contraindication to conventional therapy (such as methotrexate, sulfasalazine, leflunomide);

AND

D. Individual has had a trial (medication samples/coupons/discount cards are excluded from consideration as a trial) and inadequate response or intolerance to TWO (2) preferred biologic agents [Current preferred biologics include – Enbrel (etanercept), Humira (adalimumab)] unless the following criteria are met:

- 1. Individual has been receiving and is maintained on a stable dose of Xeljanz (tofacitinib) or Xeljanz XR (tofacitinib extended-release); **OR**
- 2. The preferred agents are not FDA-approved and do not have an accepted off-label use per the off-label policy for the prescribed indication and Xeljanz (tofacitinib) or Xeljanz XR (tofacitinib extended-release) does; **OR**
- 3. The preferred agents are not acceptable due to concomitant clinical conditions, such as but not limited to any of the following:

This policy does not apply to health plans or member categories that do not have pharmacy benefits, nor does it apply to Medicare. Note that market specific restrictions or transition-of-care benefit limitations may apply.

Market Applicability														
Market	DC	FL & FHK	FL MMA	FL LTC	GA	KS	KY	MD	NJ	NV	NY	TN	TX	WA
Applicable	X	X	NA	NA	X	NA	X	X	X	X	X	NA	NA	NA

*FHK- Florida Healthy Kids

- a. Known hypersensitivity to any active or inactive component which is not also associated with Xeljanz (tofacitinib) or Xeljanz XR (tofacitinib extended-release); **OR**
- b. Individual's age; **OR**
- c. Pregnant or planning on becoming pregnant; **OR**
- d. Serious infections or concurrent sepsis; **OR**
4. The individual has either concomitant clinical condition:
 - a. Demyelinating disease; **OR**
 - b. Heart failure with documented left ventricular dysfunction; **OR**
5. The preferred agent(s) do not have activity against a concomitant clinical condition and Xeljanz (tofacitinib) or Xeljanz XR (tofacitinib extended-release) does. Examples include but may not be limited to the following:
 - a. Concomitant Crohn's Disease: TNFi (agents FDA-approved for both indications) or Stelara are preferred; **OR**
 - b. Concomitant Ulcerative Colitis: TNFi (agents FDA-approved for both indications) are preferred.

Requests for Xeljanz (tofacitinib) may be approved based on the following criteria:

- I. Diagnosis of Ulcerative Colitis:
 - A. Individual is 18 years of age or older with moderately to severely active ulcerative colitis; **AND**
 - B. Individual has had an inadequate response to, is intolerant of, or has a contraindication to conventional therapy (such as 5-aminosalicylic acid products, systemic corticosteroids, or immunosuppressive drugs); **AND**
 - C. Individual has had a trial (medication samples/coupons/discount cards are excluded from consideration as a trial) and inadequate response or intolerance to ONE (1) preferred biologic agents [Current preferred biologics include – Humira (adalimumab), Inflectra (infliximab-dyyb), or Renflexis (infliximab-abda)] unless the following criteria are met:
 1. Individual has been receiving and is maintained on a stable dose of Xeljanz (tofacitinib); **OR**
 2. The preferred agents are not FDA-approved and do not have an accepted off-label use per the off-label policy for the prescribed indication and Xeljanz (tofacitinib) does; **OR**
 3. The preferred agents are not acceptable due to concomitant clinical conditions, such as but not limited to any of the following:

This policy does not apply to health plans or member categories that do not have pharmacy benefits, nor does it apply to Medicare. Note that market specific restrictions or transition-of-care benefit limitations may apply.

Market Applicability														
Market	DC	FL & FHK	FL MMA	FL LTC	GA	KS	KY	MD	NJ	NV	NY	TN	TX	WA
Applicable	X	X	NA	NA	X	NA	X	X	X	X	X	NA	NA	NA

*FHK- Florida Healthy Kids

- a. Known hypersensitivity to any active or inactive component which is not also associated with Xeljanz (tofacitinib); **OR**
- b. Individual's age; **OR**
- c. Pregnant or planning on becoming pregnant; **OR**
- d. Serious infections or concurrent sepsis; **OR**
4. Individual has any of the following concomitant clinical conditions:
 - a. Demyelinating disease; **OR**
 - b. Heart failure with documented left ventricular dysfunction; **OR**
 - c. Malignancy [such as but not limited to, solid or hematologic cancers and excluding superficial skin cancers (such as basal and squamous cell)]; **OR**
5. The preferred agent(s) do not have activity against a concomitant clinical condition and Xeljanz (tofacitinib) does. Examples include but may not be limited to the following:
 - a. Concomitant Psoriasis: TNFi (agents FDA-approved for both indications) or Stelara are preferred; **OR**
 - b. Concomitant Rheumatoid Arthritis: TNFi (agents FDA-approved for both indications) or Xeljanz are preferred.

Xeljanz XR (tofacitinib extended-release) may **not** be approved for the following:

- I. Individual has a diagnosis of moderate [30-59 mL/min/1.73 m² (NKF 2002, KDIGO 2012)] or severe [less than 30 mL/min/1.73 m² (NKF 2002, KDIGO 2012)] renal impairment; **OR**
- II. Individual has a diagnosis of moderate hepatic impairment (Child Pugh Class B).

Xeljanz (tofacitinib) or Xeljanz XR (tofacitinib extended-release) may **not** be approved for the following:

- I. In combination with other JAK inhibitors (such as Olumiant), biologic disease-modifying antirheumatic drugs (DMARDs) or potent immunosuppressants such as azathioprine and cyclosporine; **OR**
- II. At initiation of therapy, absolute neutrophil count (ANC) less than 1000 cells/mm³, lymphocyte count less than 500 cells/mm³, or hemoglobin less than 9 g/dL; **OR**
- III. Tuberculosis or other active serious infections or a history of recurrent infections; **OR**
- IV. Individual has not had a tuberculin skin test (TST) or a Centers for Disease Control (CDC)-recommended equivalent to evaluate for latent tuberculosis prior to initiating Xeljanz; **OR**

This policy does not apply to health plans or member categories that do not have pharmacy benefits, nor does it apply to Medicare. Note that market specific restrictions or transition-of-care benefit limitations may apply.

Market Applicability														
Market	DC	FL & FHK	FL MMA	FL LTC	GA	KS	KY	MD	NJ	NV	NY	TN	TX	WA
Applicable	X	X	NA	NA	X	NA	X	X	X	X	X	NA	NA	NA

*FHK- Florida Healthy Kids

V. Individual has severe hepatic impairment (Child Pugh class C).

Note: Xeljanz (tofacitinib), Xeljanz XR (tofacitinib extended-release) has black box warnings for serious infections and malignancy. The increased risk of developing serious infections can result in hospitalization or death. Most individuals that developed serious infections were taking concomitant immunosuppressants. Individuals should be closely monitored for the development of an infection during and after treatment with discontinuation of therapy if the individual develops a serious infection. Reported infections include: Active tuberculosis (pulmonary or extrapulmonary disease), invasive fungal infections (including cryptococcosis and pneumocystosis), and infections (bacterial, viral, or other) due to opportunistic pathogens. Individuals should be tested for latent tuberculosis prior to and during therapy. Latent tuberculosis should be treated prior to initiation of therapy. The risks and benefits of treatment with Xeljanz should be considered prior to initiating in individuals with chronic or recurrent infection. Lymphoma and other malignancies have occurred with therapy. Epstein Barr virus-associated post-transplant lymphoproliferative disorder has been observed in renal transplant individuals taking concomitant immunosuppressants.

State Specific Mandates		
State name	Date effective	Mandate details (including specific bill if applicable)
N/A	N/A	N/A

Key References:

1. Clinical Pharmacology [database online]. Tampa, FL: Gold Standard, Inc.: 2018. URL: <http://www.clinicalpharmacology.com>. Updated periodically.
2. DailyMed. Package inserts. U.S. National Library of Medicine, National Institutes of Health website. <http://dailymed.nlm.nih.gov/dailymed/about.cfm>. Accessed: August 27, 2018
3. DrugPoints® System [electronic version]. Truven Health Analytics, Greenwood Village, CO. Updated periodically.
4. Lexi-Comp ONLINE™ with AHFS™, Hudson, Ohio: Lexi-Comp, Inc.; 2018; Updated periodically.
5. Singh JA, Saag KG, Bridges SL et al. 2015 American College of Rheumatology Guideline for the treatment of rheumatoid arthritis. *Arthritis Rheum.* 2016;68:1-26.
6. American Gastroenterological Association. Identification, assessment and initial medical treatment of ulcerative colitis Clinical Care Pathway. Available at <https://gastro.org/guidelines/ibd-and-bowel-disorders>. Accessed on: August 24, 2018.
7. Kidney Disease: Improving Global Outcomes (KDIGO) CKD Work Group. KDIGO 2012 Clinical Practice Guideline for the Evaluation and Management of Chronic Kidney Disease. *Kidney Int Suppl.*2013; 3:1–150. Available from <http://kdigo.org/home/guidelines/ckd-evaluation-management/>. Accessed on: August 24, 2018.

This policy does not apply to health plans or member categories that do not have pharmacy benefits, nor does it apply to Medicare. Note that market specific restrictions or transition-of-care benefit limitations may apply.